

Algebraic expressions

1

Objectives

After completing this chapter you should be able to:

- Multiply and divide integer powers → pages 2–3
- Expand a single term over brackets and collect like terms → pages 3–4
- Expand the product of two or three expressions → pages 4–6
- Factorise linear, quadratic and simple cubic expressions → pages 6–9
- Know and use the laws of indices → pages 9–11
- Simplify and use the rules of surds → pages 12–13
- Rationalise denominators → pages 13–16

Prior knowledge check

1 Simplify:

a $4m^2n + 5mn^2 - 2m^2n + mn^2 - 3mn^2$

b $3x^2 - 5x + 2 + 3x^2 - 7x - 12$

← GCSE Mathematics

2 Write as a single power of 2:

a $2^5 \times 2^3$

b $2^6 \div 2^2$

c $(2^3)^2$

← GCSE Mathematics

3 Expand:

a $3(x + 4)$

b $5(2 - 3x)$

c $6(2x - 5y)$

← GCSE Mathematics

4 Write down the highest common factor of:

a 24 and 16

b $6x$ and $8x^2$

c $4xy^2$ and $3xy$

← GCSE Mathematics

5 Simplify:

a $\frac{10x}{5}$

b $\frac{20x}{2}$

c $\frac{40x}{24}$

← GCSE Mathematics

Computer scientists use indices to describe very large numbers. A quantum computer with 1000 qubits (quantum bits) can consider 2^{1000} values simultaneously. This is greater than the number of particles in the observable universe.

1.1 Index laws

■ You can use the laws of indices to simplify powers of the same base.

- $a^m \times a^n = a^{m+n}$
- $a^m \div a^n = a^{m-n}$
- $(a^m)^n = a^{mn}$
- $(ab)^n = a^n b^n$

Notation

x^5 This is the **base**.
This is the **index, power or exponent**.

Example 1

Simplify these expressions:

a $x^2 \times x^5$ b $2r^2 \times 3r^3$ c $\frac{b^7}{b^4}$ d $6x^5 \div 3x^3$ e $(a^3)^2 \times 2a^2$ f $(3x^2)^3 \div x^4$

a $x^2 \times x^5 = x^{2+5} = x^7$

Use the rule $a^m \times a^n = a^{m+n}$ to simplify the index.

b $2r^2 \times 3r^3 = 2 \times 3 \times r^2 \times r^3$
 $= 6 \times r^{2+3} = 6r^5$

Rewrite the expression with the numbers together and the r terms together.

c $\frac{b^7}{b^4} = b^{7-4} = b^3$

$2 \times 3 = 6$

$r^2 \times r^3 = r^{2+3}$

d $6x^5 \div 3x^3 = \frac{6}{3} \times \frac{x^5}{x^3}$
 $= 2 \times x^2 = 2x^2$

Use the rule $a^m \div a^n = a^{m-n}$ to simplify the index.

e $(a^3)^2 \times 2a^2 = a^6 \times 2a^2$
 $= 2 \times a^6 \times a^2 = 2a^8$

$x^5 \div x^3 = x^{5-3} = x^2$

Use the rule $(a^m)^n = a^{mn}$ to simplify the index.

f $\frac{(3x^2)^3}{x^4} = 3^3 \times \frac{(x^2)^3}{x^4}$
 $= 27 \times \frac{x^6}{x^4} = 27x^2$

$a^6 \times a^2 = a^{6+2} = a^8$

Use the rule $(ab)^n = a^n b^n$ to simplify the numerator.

$(x^2)^3 = x^{2 \times 3} = x^6$

$\frac{x^6}{x^4} = x^{6-4} = x^2$

Example 2

Expand these expressions and simplify if possible:

a $-3x(7x-4)$

b $y^2(3-2y^3)$

c $4x(3x-2x^2+5x^3)$

d $2x(5x+3)-5(2x+3)$

Watch out

A minus sign outside brackets changes the sign of every term inside the brackets.

$$a \quad -3x(7x - 4) = -21x^2 + 12x$$

$$b \quad y^2(3 - 2y^3) = 3y^2 - 2y^5$$

$$c \quad 4x(3x - 2x^2 + 5x^3) \\ = 12x^2 - 8x^3 + 20x^4$$

$$d \quad 2x(5x + 3) - 5(2x + 3) \\ = 10x^2 + 6x - 10x - 15 \\ = 10x^2 - 4x - 15$$

$$-3x \times 7x = -21x^{1+1} = -21x^2$$

$$-3x \times (-4) = +12x$$

$$y^2 \times (-2y^3) = -2y^{2+3} = -2y^5$$

Remember a minus sign outside the brackets changes the signs within the brackets.

Simplify $6x - 10x$ to give $-4x$.

Example 3

Simplify these expressions:

$$a \quad \frac{x^7 + x^4}{x^3} \quad b \quad \frac{3x^2 - 6x^5}{2x} \quad c \quad \frac{20x^7 + 15x^3}{5x^2}$$

$$a \quad \frac{x^7 + x^4}{x^3} = \frac{x^7}{x^3} + \frac{x^4}{x^3} \\ = x^{7-3} + x^{4-3} = x^4 + x$$

$$b \quad \frac{3x^2 - 6x^5}{2x} = \frac{3x^2}{2x} - \frac{6x^5}{2x} \\ = \frac{3}{2}x^{2-1} - 3x^{5-1} = \frac{3x}{2} - 3x^4$$

$$c \quad \frac{20x^7 + 15x^3}{5x^2} = \frac{20x^7}{5x^2} + \frac{15x^3}{5x^2} \\ = 4x^{7-2} + 3x^{3-2} = 4x^5 + 3x$$

Divide each term of the numerator by x^3 .

x^1 is the same as x .

Divide each term of the numerator by $2x$.

Simplify each fraction:

$$\frac{3x^2}{2x} = \frac{3}{2} \times \frac{x^2}{x} = \frac{3}{2} \times x^{2-1}$$

$$-\frac{6x^5}{2x} = -\frac{6}{2} \times \frac{x^5}{x} = -3 \times x^{5-1}$$

Divide each term of the numerator by $5x^2$.



Exercise 1A

1 Simplify these expressions:

$$a \quad x^3 \times x^4$$

$$b \quad 2x^3 \times 3x^2$$

$$c \quad \frac{k^3}{k^2}$$

$$d \quad \frac{4p^3}{2p}$$

$$e \quad \frac{3x^3}{3x^2}$$

$$f \quad (y^2)^5$$

$$g \quad 10x^5 \div 2x^3$$

$$h \quad (p^3)^2 \div p^4$$

$$i \quad (2a^3)^2 \div 2a^3$$

$$j \quad 8p^4 \div 4p^3$$

$$k \quad 2a^4 \times 3a^5$$

$$l \quad \frac{21a^3b^7}{7ab^4}$$

$$m \quad 9x^2 \times 3(x^2)^3$$

$$n \quad 3x^3 \times 2x^2 \times 4x^6$$

$$o \quad 7a^4 \times (3a^4)^2$$

$$p \quad (4y^3)^3 \div 2y^3$$

$$q \quad 2a^3 \div 3a^2 \times 6a^5$$

$$r \quad 3a^4 \times 2a^5 \times a^3$$

2 Expand and simplify if possible:

- | | | |
|------------------------------------|--|---------------------------------|
| a $9(x - 2)$ | b $x(x + 9)$ | c $-3y(4 - 3y)$ |
| d $x(y + 5)$ | e $-x(3x + 5)$ | f $-5x(4x + 1)$ |
| g $(4x + 5)x$ | h $-3y(5 - 2y^2)$ | i $-2x(5x - 4)$ |
| j $(3x - 5)x^2$ | k $3(x + 2) + (x - 7)$ | l $5x - 6 - (3x - 2)$ |
| m $4(c + 3d^2) - 3(2c + d^2)$ | n $(r^2 + 3t^2 + 9) - (2r^2 + 3t^2 - 4)$ | |
| o $x(3x^2 - 2x + 5)$ | p $7y^2(2 - 5y + 3y^2)$ | q $-2y^2(5 - 7y + 3y^2)$ |
| r $7(x - 2) + 3(x + 4) - 6(x - 2)$ | | s $5x - 3(4 - 2x) + 6$ |
| t $3x^2 - x(3 - 4x) + 7$ | u $4x(x + 3) - 2x(3x - 7)$ | v $3x^2(2x + 1) - 5x^2(3x - 4)$ |

3 Simplify these fractions:

- | | | |
|-----------------------------|----------------------------|----------------------------|
| a $\frac{6x^4 + 10x^6}{2x}$ | b $\frac{3x^5 - x^7}{x}$ | c $\frac{2x^4 - 4x^2}{4x}$ |
| d $\frac{8x^3 + 5x}{2x}$ | e $\frac{7x^7 + 5x^2}{5x}$ | f $\frac{9x^5 - 5x^3}{3x}$ |

1.2 Expanding brackets

To find the **product** of two expressions you **multiply** each term in one expression by each term in the other expression.

Multiplying each of the 2 terms in the first expression by each of the 3 terms in the second expression gives $2 \times 3 = 6$ terms.

$$\begin{aligned}
 (x + 5)(4x - 2y + 3) &= x(4x - 2y + 3) + 5(4x - 2y + 3) \\
 &= 4x^2 - 2xy + 3x + 20x - 10y + 15 \\
 &= 4x^2 - 2xy + 23x - 10y + 15
 \end{aligned}$$

Simplify your answer by collecting like terms.

Example 4

Expand these expressions and simplify if possible:

- a $(x + 5)(x + 2)$ b $(x - 2y)(x^2 + 1)$ c $(x - y)^2$ d $(x + y)(3x - 2y - 4)$

a $(x + 5)(x + 2)$

$$\begin{aligned}
 &= x^2 + 2x + 5x + 10 \\
 &= x^2 + 7x + 10
 \end{aligned}$$

Multiply x by $(x + 2)$ and then multiply 5 by $(x + 2)$.

Simplify your answer by collecting like terms.

b $(x - 2y)(x^2 + 1)$

$$\begin{aligned}
 &= x^3 + x - 2x^2y - 2y
 \end{aligned}$$

$$-2y \times x^2 = -2x^2y$$

There are no like terms to collect.

$$\begin{aligned}
 \text{c } (x - y)^2 &= (x - y)(x - y) \\
 &= x^2 - \underline{xy} - \underline{xy} + y^2 \\
 &= x^2 - 2xy + y^2
 \end{aligned}$$

$(x - y)^2$ means $(x - y)$ multiplied by itself.

$$-xy - xy = -2xy$$

$$\begin{aligned}
 \text{d } (x + y)(3x - 2y - 4) &= x(3x - 2y - 4) + y(3x - 2y - 4) \\
 &= 3x^2 - 2xy - 4x + 3xy - 2y^2 - 4y \\
 &= 3x^2 + xy - 4x - 2y^2 - 4y
 \end{aligned}$$

Multiply x by $(3x - 2y - 4)$ and then multiply y by $(3x - 2y - 4)$.

Example 5

Expand these expressions and simplify if possible:

a $x(2x + 3)(x - 7)$

b $x(5x - 3y)(2x - y + 4)$

c $(x - 4)(x + 3)(x + 1)$

$$\begin{aligned}
 \text{a } x(2x + 3)(x - 7) &= (2x^2 + 3x)(x - 7) \\
 &= 2x^3 - 14x^2 + 3x^2 - 21x \\
 &= 2x^3 - 11x^2 - 21x
 \end{aligned}$$

Start by expanding one pair of brackets:
 $x(2x + 3) = 2x^2 + 3x$

You could also have expanded the second pair of brackets first: $(2x + 3)(x - 7) = 2x^2 - 11x - 21$
 Then multiply by x .

$$\begin{aligned}
 \text{b } x(5x - 3y)(2x - y + 4) &= (5x^2 - 3xy)(2x - y + 4) \\
 &= 5x^2(2x - y + 4) - 3xy(2x - y + 4) \\
 &= 10x^3 - 5x^2y + 20x^2 - 6x^2y + 3xy^2 - 12xy \\
 &= 10x^3 - 11x^2y + 20x^2 + 3xy^2 - 12xy
 \end{aligned}$$

Be careful with minus signs. You need to change every sign in the second pair of brackets when you multiply it out.

Choose one pair of brackets to expand first, for example:

$$\begin{aligned}
 (x - 4)(x + 3) &= x^2 + 3x - 4x - 12 \\
 &= x^2 - x - 12
 \end{aligned}$$

$$\begin{aligned}
 \text{c } (x - 4)(x + 3)(x + 1) &= (x^2 - x - 12)(x + 1) \\
 &= x^2(x + 1) - x(x + 1) - 12(x + 1) \\
 &= x^3 + x^2 - x^2 - x - 12x - 12 \\
 &= x^3 - 13x - 12
 \end{aligned}$$

You multiplied together three linear terms, so the final answer contains an x^3 term.



Exercise 1B

1 Expand and simplify if possible:

a $(x + 4)(x + 7)$

b $(x - 3)(x + 2)$

c $(x - 2)^2$

d $(x - y)(2x + 3)$

e $(x + 3y)(4x - y)$

f $(2x - 4y)(3x + y)$

g $(2x - 3)(x - 4)$

h $(3x + 2y)^2$

i $(2x + 8y)(2x + 3)$

j $(x + 5)(2x + 3y - 5)$

k $(x - 1)(3x - 4y - 5)$

l $(x - 4y)(2x + y + 5)$

m $(x + 2y - 1)(x + 3)$

n $(2x + 2y + 3)(x + 6)$

o $(4 - y)(4y - x + 3)$

p $(4y + 5)(3x - y + 2)$

q $(5y - 2x + 3)(x - 4)$

r $(4y - x - 2)(5 - y)$

2 Expand and simplify if possible:

a $5(x+1)(x-4)$

b $7(x-2)(2x+5)$

c $3(x-3)(x-3)$

d $x(x-y)(x+y)$

e $x(2x+y)(3x+4)$

f $y(x-5)(x+1)$

g $y(3x-2y)(4x+2)$

h $y(7-x)(2x-5)$

i $x(2x+y)(5x-2)$

j $x(x+2)(x+3y-4)$

k $y(2x+y-1)(x+5)$

l $y(3x+2y-3)(2x+1)$

m $x(2x+3)(x+y-5)$

n $2x(3x-1)(4x-y-3)$

o $3x(x-2y)(2x+3y+5)$

p $(x+3)(x+2)(x+1)$

q $(x+2)(x-4)(x+3)$

r $(x+3)(x-1)(x-5)$

s $(x-5)(x-4)(x-3)$

t $(2x+1)(x-2)(x+1)$


u $(2x+3)(3x-1)(x+2)$

v $(3x-2)(2x+1)(3x-2)$

w $(x+y)(x-y)(x-1)$


x $(2x-3y)^3$

- (P) 3 The diagram shows a rectangle with a square cut out. The rectangle has length $3x - y + 4$ and width $x + 7$. The square has length $x - 2$. Find an expanded and simplified expression for the shaded area.



Problem-solving

Use the same strategy as you would use if the lengths were given as numbers:



- (P) 4 A cuboid has dimensions $x + 2$ cm, $2x - 1$ cm and $2x + 3$ cm. Show that the volume of the cuboid is $4x^3 + 12x^2 + 5x - 6$ cm³.
- (E/P) 5 Given that $(2x + 5y)(3x - y)(2x + y) = ax^3 + bx^2y + cxy^2 + dy^3$, where a , b , c and d are constants, find the values of a , b , c and d . (2 marks)

Challenge

Expand and simplify $(x + y)^4$.

Links

You can use the binomial expansion to expand expressions like $(x + y)^4$ quickly. → Section 8.3

1.3 Factorising

You can write expressions as a **product of their factors**.

- Factorising is the opposite of expanding brackets.

Expanding brackets

$$4x(2x + y) = 8x^2 + 4xy$$

$$(x + 5)^3 = x^3 + 15x^2 + 75x + 125$$

$$(x + 2y)(x - 5y) = x^2 - 3xy - 10y^2$$

Factorising

Example 6

Factorise these expressions completely:

a $3x + 9$

b $x^2 - 5x$

c $8x^2 + 20x$

d $9x^2y + 15xy^2$

e $3x^2 - 9xy$

a $3x + 9 = 3(x + 3)$

3 is a common factor of $3x$ and 9.

b $x^2 - 5x = x(x - 5)$

 x is a common factor of x^2 and $-5x$.

c $8x^2 + 20x = 4x(2x + 5)$

4 and x are common factors of $8x^2$ and $20x$.
So take $4x$ outside the brackets.

d $9x^2y + 15xy^2 = 3xy(3x + 5y)$

3, x and y are common factors of $9x^2y$ and $15xy^2$.
So take $3xy$ outside the brackets.

e $3x^2 - 9xy = 3x(x - 3y)$

 x and $-3y$ have no common factors so this expression is completely factorised.

- A quadratic expression has the form $ax^2 + bx + c$ where a , b and c are real numbers and $a \neq 0$.

Notation Real numbers are all the positive and negative numbers, or zero, including fractions and surds.

To factorise a quadratic expression:

- Find two factors of ac that add up to b
- Rewrite the b term as a sum of these two factors
- Factorise each pair of terms
- Take out the common factor

For the expression $2x^2 + 5x - 3$, $ac = -6 = -1 \times 6$
and $-1 + 6 = 5 = b$.

$2x^2 - x + 6x - 3$

$= x(2x - 1) + 3(2x - 1)$

$= (x + 3)(2x - 1)$

■ $x^2 - y^2 = (x + y)(x - y)$

Notation An expression in the form $x^2 - y^2$ is called the **difference** of two squares.

Example 7

Factorise:

a $x^2 - 5x - 6$

b $x^2 + 6x + 8$

c $6x^2 - 11x - 10$

d $x^2 - 25$

e $4x^2 - 9y^2$

a $x^2 - 5x - 6$

$ac = -6$ and $b = -5$

So $x^2 - 5x - 6 = x^2 + x - 6x - 6$

$= x(x + 1) - 6(x + 1)$

$= (x + 1)(x - 6)$

Here $a = 1$, $b = -5$ and $c = -6$.① Work out the two factors of $ac = -6$ which add to give you $b = -5$. $-6 + 1 = -5$ ② Rewrite the b term using these two factors.

③ Factorise first two terms and last two terms.

④ $x + 1$ is a factor of both terms, so take that outside the brackets. This is now completely factorised.

b $x^2 + 6x + 8$

$$= x^2 + 2x + 4x + 8$$

$$= x(x + 2) + 4(x + 2)$$

$$= (x + 2)(x + 4)$$

$$ac = 8 \text{ and } 2 + 4 = 6 = b.$$

Factorise.

c $6x^2 - 11x - 10$

$$= 6x^2 - 15x + 4x - 10$$

$$= 3x(2x - 5) + 2(2x - 5)$$

$$= (2x - 5)(3x + 2)$$

$$ac = -60 \text{ and } 4 - 15 = -11 = b.$$

Factorise.

d $x^2 - 25$

$$= x^2 - 5^2$$

$$= (x + 5)(x - 5)$$

This is the difference of two squares as the two terms are x^2 and 5^2 .

The two x terms, $5x$ and $-5x$, cancel each other out.

e $4x^2 - 9y^2$

$$= 2^2x^2 - 3^2y^2$$

$$= (2x + 3y)(2x - 3y)$$

$$\text{This is the same as } (2x)^2 - (3y)^2.$$

Example 8

Factorise completely:

a $x^3 - 2x^2$ **b** $x^3 - 25x$ **c** $x^3 + 3x^2 - 10x$

a $x^3 - 2x^2 = x^2(x - 2)$

You can't factorise this any further.

b $x^3 - 25x = x(x^2 - 25)$

$$= x(x^2 - 5^2)$$

$$= x(x + 5)(x - 5)$$

x is a common factor of x^3 and $-25x$.
So take x outside the brackets.

c $x^3 + 3x^2 - 10x = x(x^2 + 3x - 10)$

$x^2 - 25$ is the difference of two squares.

$$= x(x + 5)(x - 2)$$

Write the expression as a product of x and a quadratic factor.

Factorise the quadratic to get three linear factors.

Exercise 1C

1 Factorise these expressions completely:

a $4x + 8$

b $6x - 24$

c $20x + 15$

d $2x^2 + 4$

e $4x^2 + 20$

f $6x^2 - 18x$

g $x^2 - 7x$

h $2x^2 + 4x$

i $3x^2 - x$

j $6x^2 - 2x$

k $10y^2 - 5y$

l $35x^2 - 28x$

m $x^2 + 2x$

n $3y^2 + 2y$

o $4x^2 + 12x$

p $5y^2 - 20y$

q $9xy^2 + 12x^2y$

r $6ab - 2ab^2$

s $5x^2 - 25xy$

t $12x^2y + 8xy^2$

u $15y - 20yz^2$

v $12x^2 - 30$

w $xy^2 - x^2y$

x $12y^2 - 4yx$

2 Factorise:

a $x^2 + 4x$

d $x^2 + 8x + 12$

g $x^2 + 5x + 6$

j $x^2 + x - 20$

m $5x^2 - 16x + 3$

o $2x^2 + 7x - 15$

q $x^2 - 4$

s $4x^2 - 25$

v $2x^2 - 50$

b $2x^2 + 6x$

e $x^2 + 3x - 40$

h $x^2 - 2x - 24$

k $2x^2 + 5x + 2$

n $6x^2 - 8x - 8$

p $2x^4 + 14x^2 + 24$

r $x^2 - 49$

t $9x^2 - 25y^2$

w $6x^2 - 10x + 4$

c $x^2 + 11x + 24$

f $x^2 - 8x + 12$

i $x^2 - 3x - 10$

l $3x^2 + 10x - 8$

Hint For part **n**, take 2 out as a common factor first. For part **p**, let $y = x^2$.

u $36x^2 - 4$

x $15x^2 + 42x - 9$

3 Factorise completely:

a $x^3 + 2x$

d $x^3 - 9x$

g $x^3 - 7x^2 + 6x$

j $2x^3 + 13x^2 + 15x$

b $x^3 - x^2 + x$

e $x^3 - x^2 - 12x$

h $x^3 - 64x$

k $x^3 - 4x$

c $x^3 - 5x$

f $x^3 + 11x^2 + 30x$

i $2x^3 - 5x^2 - 3x$

l $3x^3 + 27x^2 + 60x$

E/P 4 Factorise completely $x^4 - y^4$. (2 marks)

Problem-solving

Watch out for terms that can be written as a function of a function: $x^4 = (x^2)^2$

E 5 Factorise completely $6x^3 + 7x^2 - 5x$. (2 marks)

Challenge

Write $4x^4 - 13x^2 + 9$ as the product of four linear factors.

1.4 Negative and fractional indices

Indices can be negative numbers or fractions.

$$x^{\frac{1}{2}} \times x^{\frac{1}{2}} = x^{\frac{1}{2} + \frac{1}{2}} = x^1 = x,$$

similarly $\underbrace{x^{\frac{1}{n}} \times x^{\frac{1}{n}} \times \dots \times x^{\frac{1}{n}}}_{n \text{ terms}} = x^{\frac{1}{n} + \frac{1}{n} + \dots + \frac{1}{n}} = x^1 = x$

Notation **Rational**
numbers are those that
can be written as $\frac{a}{b}$ where
 a and b are integers.

■ You can use the laws of indices with any rational power.

- $a^{\frac{1}{m}} = \sqrt[m]{a}$
- $a^{\frac{n}{m}} = \sqrt[m]{a^n}$
- $a^{-m} = \frac{1}{a^m}$
- $a^0 = 1$

Notation $a^{\frac{1}{2}} = \sqrt{a}$ is the
positive square root of a .
For example $9^{\frac{1}{2}} = \sqrt{9} = 3$
but $9^{\frac{1}{2}} \neq -3$.

Example 9

Simplify:

a $\frac{x^3}{x^{-3}}$

b $x^{\frac{1}{2}} \times x^{\frac{3}{2}}$

c $(x^3)^{\frac{2}{3}}$

d $2x^{1.5} \div 4x^{-0.25}$

e $\sqrt[3]{125x^6}$

f $\frac{2x^2 - x}{x^5}$

a $\frac{x^3}{x^{-3}} = x^{3 - (-3)} = x^6$

Use the rule $a^m \div a^n = a^{m-n}$.

b $x^{\frac{1}{2}} \times x^{\frac{3}{2}} = x^{\frac{1}{2} + \frac{3}{2}} = x^2$

This could also be written as \sqrt{x} .
Use the rule $a^m \times a^n = a^{m+n}$.

c $(x^3)^{\frac{2}{3}} = x^3 \times \frac{2}{3} = x^2$

Use the rule $(a^m)^n = a^{mn}$.

d $2x^{1.5} \div 4x^{-0.25} = \frac{1}{2}x^{1.5 - (-0.25)} = \frac{1}{2}x^{1.75}$

Use the rule $a^m \div a^n = a^{m-n}$.
 $1.5 - (-0.25) = 1.75$

e $\sqrt[3]{125x^6} = (125x^6)^{\frac{1}{3}}$
 $= (125)^{\frac{1}{3}}(x^6)^{\frac{1}{3}} = \sqrt[3]{125}(x^{6 \times \frac{1}{3}}) = 5x^2$

Using $a^{\frac{1}{m}} = \sqrt[m]{a}$.

f $\frac{2x^2 - x}{x^5} = \frac{2x^2}{x^5} - \frac{x}{x^5}$
 $= 2 \times x^{2-5} - x^{1-5} = 2x^{-3} - x^{-4}$
 $= \frac{2}{x^3} - \frac{1}{x^4}$

Divide each term of the numerator by x^5 .Using $a^{-m} = \frac{1}{a^m}$ **Example 10**

Evaluate:

a $9^{\frac{1}{2}}$

b $64^{\frac{1}{3}}$

c $49^{\frac{3}{2}}$

d $25^{-\frac{3}{2}}$

a $9^{\frac{1}{2}} = \sqrt{9} = 3$

Using $a^{\frac{1}{m}} = \sqrt[m]{a}$. $9^{\frac{1}{2}} = \sqrt{9}$

b $64^{\frac{1}{3}} = \sqrt[3]{64} = 4$

This means the cube root of 64.

c $49^{\frac{3}{2}} = (\sqrt{49})^3$
 $7^3 = 343$

Using $a^{\frac{n}{m}} = \sqrt[m]{a^n}$.
This means the square root of 49, cubed.

d $25^{-\frac{3}{2}} = \frac{1}{25^{\frac{3}{2}}} = \frac{1}{(\sqrt{25})^3}$
 $= \frac{1}{5^3} = \frac{1}{125}$

Using $a^{-m} = \frac{1}{a^m}$ **Online** Use your calculator to enter negative and fractional powers.

Example 11

Given that $y = \frac{1}{16}x^2$ express each of the following in the form kx^n , where k and n are constants.

a $y^{\frac{1}{2}}$

b $4y^{-1}$

$$\begin{aligned} \text{a } y^{\frac{1}{2}} &= \left(\frac{1}{16}x^2\right)^{\frac{1}{2}} \\ &= \frac{1}{\sqrt{16}}x^{2 \times \frac{1}{2}} = \frac{x}{4} \end{aligned}$$

$$\begin{aligned} \text{b } 4y^{-1} &= 4\left(\frac{1}{16}x^2\right)^{-1} \\ &= 4\left(\frac{1}{16}\right)^{-1}x^{2 \times (-1)} = 4 \times 16x^{-2} \\ &= 64x^{-2} \end{aligned}$$

Substitute $y = \frac{1}{16}x^2$ into $y^{\frac{1}{2}}$.

$$\left(\frac{1}{16}\right)^{\frac{1}{2}} = \frac{1}{\sqrt{16}} \text{ and } (x^2)^{\frac{1}{2}} = x^{2 \times \frac{1}{2}}$$

$$\left(\frac{1}{16}\right)^{-1} = 16 \text{ and } x^{2 \times (-1)} = x^{-2}$$

Problem-solving

Check that your answers are in the correct form. If k and n are constants they could be positive or negative, and they could be integers, fractions or surds.

Exercise 1D

1 Simplify:

a $x^3 \div x^{-2}$

b $x^5 \div x^7$

c $x^{\frac{3}{2}} \times x^{\frac{5}{2}}$

d $(x^2)^{\frac{3}{2}}$

e $(x^3)^{\frac{5}{3}}$

f $3x^{0.5} \times 4x^{-0.5}$

g $9x^{\frac{2}{3}} \div 3x^{\frac{1}{6}}$

h $5x^{\frac{7}{2}} \div x^{\frac{3}{2}}$

i $3x^4 \times 2x^{-5}$

j $\sqrt{x} \times \sqrt[3]{x}$

k $(\sqrt{x})^3 \times (\sqrt[3]{x})^4$

l $\frac{(\sqrt[3]{x})^2}{\sqrt{x}}$

2 Evaluate:

a $25^{\frac{1}{2}}$

b $81^{\frac{3}{2}}$

c $27^{\frac{1}{3}}$

d 4^{-2}

e $9^{-\frac{1}{2}}$

f $(-5)^{-3}$

g $\left(\frac{3}{4}\right)^0$

h $1296^{\frac{3}{4}}$

i $\left(\frac{25}{16}\right)^{\frac{3}{2}}$

j $\left(\frac{27}{8}\right)^{\frac{2}{3}}$

k $\left(\frac{6}{5}\right)^{-1}$

l $\left(\frac{343}{512}\right)^{-\frac{2}{3}}$

3 Simplify:

a $(64x^{10})^{\frac{1}{2}}$

b $\frac{5x^3 - 2x^2}{x^5}$

c $(125x^{12})^{\frac{1}{3}}$

d $\frac{x + 4x^3}{x^3}$

e $\frac{2x + x^2}{x^4}$

f $\left(\frac{4}{9}x^4\right)^{\frac{3}{2}}$

g $\frac{9x^2 - 15x^5}{3x^3}$

h $\frac{5x + 3x^2}{15x^3}$

(E) 4 a Find the value of $81^{\frac{1}{4}}$.

(1 mark)

b Simplify $x(2x^{-\frac{1}{3}})^4$.

(2 marks)

(E) 5 Given that $y = \frac{1}{8}x^3$ express each of the following in the form kx^n , where k and n are constants.

a $y^{\frac{1}{3}}$

(2 marks)

b $\frac{1}{2}y^{-2}$

(2 marks)

1.5 Surds

If n is an integer that is **not** a square number, then any multiple of \sqrt{n} is called a surd.

Examples of surds are $\sqrt{2}$, $\sqrt{19}$ and $5\sqrt{2}$.

Surds are examples of **irrational numbers**.

The decimal expansion of a surd is never-ending and never repeats, for example $\sqrt{2} = 1.414213562\dots$

Notation Irrational numbers cannot be written in the form $\frac{a}{b}$ where a and b are integers. Surds are examples of **irrational numbers**.

You can use surds to write exact answers to calculations.

■ You can manipulate surds using these rules:

- $\sqrt{ab} = \sqrt{a} \times \sqrt{b}$
- $\sqrt{\frac{a}{b}} = \frac{\sqrt{a}}{\sqrt{b}}$

Example 12

Simplify:

a $\sqrt{12}$

b $\frac{\sqrt{20}}{2}$

c $5\sqrt{6} - 2\sqrt{24} + \sqrt{294}$

a $\sqrt{12} = \sqrt{4 \times 3}$

$= \sqrt{4} \times \sqrt{3} = 2\sqrt{3}$

b $\frac{\sqrt{20}}{2} = \frac{\sqrt{4 \times 5}}{2}$

$= \frac{2 \times \sqrt{5}}{2} = \sqrt{5}$

c $5\sqrt{6} - 2\sqrt{24} + \sqrt{294}$

$= 5\sqrt{6} - 2\sqrt{6}\sqrt{4} + \sqrt{6} \times \sqrt{49}$

$= \sqrt{6}(5 - 2\sqrt{4} + \sqrt{49})$

$= \sqrt{6}(5 - 2 \times 2 + 7)$

$= \sqrt{6}(8)$

$= 8\sqrt{6}$

Look for a factor of 12 that is a square number. Use the rule $\sqrt{ab} = \sqrt{a} \times \sqrt{b}$. $\sqrt{4} = 2$

$\sqrt{20} = \sqrt{4} \times \sqrt{5}$

$\sqrt{4} = 2$

Cancel by 2.

$\sqrt{6}$ is a common factor.

Work out the square roots $\sqrt{4}$ and $\sqrt{49}$.

$5 - 4 + 7 = 8$

Example 13

Expand and simplify if possible:

a $\sqrt{2}(5 - \sqrt{3})$

b $(2 - \sqrt{3})(5 + \sqrt{3})$

a $\sqrt{2}(5 - \sqrt{3})$

$= 5\sqrt{2} - \sqrt{2}\sqrt{3}$

$= 5\sqrt{2} - \sqrt{6}$

$\sqrt{2} \times 5 - \sqrt{2} \times \sqrt{3}$

Using $\sqrt{a} \times \sqrt{b} = \sqrt{ab}$

b $(2 - \sqrt{3})(5 + \sqrt{3})$

$= 2(5 + \sqrt{3}) - \sqrt{3}(5 + \sqrt{3})$

$= 10 + 2\sqrt{3} - 5\sqrt{3} - \sqrt{9}$

$= 7 - 3\sqrt{3}$

Expand the brackets completely before you simplify.

Collect like terms: $2\sqrt{3} - 5\sqrt{3} = -3\sqrt{3}$

Simplify any roots if possible: $\sqrt{9} = 3$

**Exercise 1E****1** Do not use your calculator for this exercise. Simplify:

a $\sqrt{28}$

b $\sqrt{72}$

c $\sqrt{50}$

d $\sqrt{32}$

e $\sqrt{90}$

f $\frac{\sqrt{12}}{2}$

g $\frac{\sqrt{27}}{3}$

h $\sqrt{20} + \sqrt{80}$

i $\sqrt{200} + \sqrt{18} - \sqrt{72}$

j $\sqrt{175} + \sqrt{63} + 2\sqrt{28}$

k $\sqrt{28} - 2\sqrt{63} + \sqrt{7}$

l $\sqrt{80} - 2\sqrt{20} + 3\sqrt{45}$

m $3\sqrt{80} - 2\sqrt{20} + 5\sqrt{45}$

n $\frac{\sqrt{44}}{\sqrt{11}}$

o $\sqrt{12} + 3\sqrt{48} + \sqrt{75}$

2 Expand and simplify if possible:

a $\sqrt{3}(2 + \sqrt{3})$

b $\sqrt{5}(3 - \sqrt{3})$

c $\sqrt{2}(4 - \sqrt{5})$

d $(2 - \sqrt{2})(3 + \sqrt{5})$

e $(2 - \sqrt{3})(3 - \sqrt{7})$

f $(4 + \sqrt{5})(2 + \sqrt{5})$

g $(5 - \sqrt{3})(1 - \sqrt{3})$

h $(4 + \sqrt{3})(2 - \sqrt{3})$

i $(7 - \sqrt{11})(2 + \sqrt{11})$

E 3 Simplify $\sqrt{75} - \sqrt{12}$ giving your answer in the form $a\sqrt{3}$, where a is an integer. **(2 marks)****1.6 Rationalising denominators**If a fraction has a surd in the denominator, it is sometimes useful to **rearrange** it so that the denominator is a **rational** number. This is called rationalising the denominator.■ **The rules to rationalise denominators are:**

- For fractions in the form $\frac{1}{\sqrt{a}}$, multiply the numerator and denominator by \sqrt{a} .
- For fractions in the form $\frac{1}{a + \sqrt{b}}$, multiply the numerator and denominator by $a - \sqrt{b}$.
- For fractions in the form $\frac{1}{a - \sqrt{b}}$, multiply the numerator and denominator by $a + \sqrt{b}$.

Example 14

Rationalise the denominator of:

a $\frac{1}{\sqrt{3}}$

b $\frac{1}{3 + \sqrt{2}}$

c $\frac{\sqrt{5} + \sqrt{2}}{\sqrt{5} - \sqrt{2}}$

d $\frac{1}{(1 - \sqrt{3})^2}$

$$\begin{aligned} \text{a } \frac{1}{\sqrt{3}} &= \frac{1 \times \sqrt{3}}{\sqrt{3} \times \sqrt{3}} \\ &= \frac{\sqrt{3}}{3} \end{aligned}$$

Multiply the numerator and denominator by $\sqrt{3}$.

$\sqrt{3} \times \sqrt{3} = (\sqrt{3})^2 = 3$

$$\begin{aligned} \text{b } \frac{1}{3 + \sqrt{2}} &= \frac{1 \times (3 - \sqrt{2})}{(3 + \sqrt{2})(3 - \sqrt{2})} \\ &= \frac{3 - \sqrt{2}}{9 - 3\sqrt{2} + 3\sqrt{2} - 2} \\ &= \frac{3 - \sqrt{2}}{7} \end{aligned}$$

Multiply numerator and denominator by $(3 - \sqrt{2})$.

$\sqrt{2} \times \sqrt{2} = 2$

$9 - 2 = 7, -3\sqrt{2} + 3\sqrt{2} = 0$

$$\begin{aligned} \text{c } \frac{\sqrt{5} + \sqrt{2}}{\sqrt{5} - \sqrt{2}} &= \frac{(\sqrt{5} + \sqrt{2})(\sqrt{5} + \sqrt{2})}{(\sqrt{5} - \sqrt{2})(\sqrt{5} + \sqrt{2})} \\ &= \frac{5 + \sqrt{5}\sqrt{2} + \sqrt{2}\sqrt{5} + 2}{5 - 2} \\ &= \frac{7 + 2\sqrt{10}}{3} \end{aligned}$$

Multiply numerator and denominator by $\sqrt{5} + \sqrt{2}$. $-\sqrt{2}\sqrt{5}$ and $\sqrt{5}\sqrt{2}$ cancel each other out.

$\sqrt{5}\sqrt{2} = \sqrt{10}$

$$\text{d } \frac{1}{(1 - \sqrt{3})^2} = \frac{1}{(1 - \sqrt{3})(1 - \sqrt{3})}$$

Expand the brackets.

$$= \frac{1}{1 - \sqrt{3} - \sqrt{3} + \sqrt{9}}$$

Simplify and collect like terms. $\sqrt{9} = 3$

$$= \frac{1}{4 - 2\sqrt{3}}$$

$$= \frac{1 \times (4 + 2\sqrt{3})}{(4 - 2\sqrt{3})(4 + 2\sqrt{3})}$$

Multiply the numerator and denominator by $4 + 2\sqrt{3}$.

$$= \frac{4 + 2\sqrt{3}}{16 + 8\sqrt{3} - 8\sqrt{3} - 12}$$

$\sqrt{3} \times \sqrt{3} = 3$

$$= \frac{4 + 2\sqrt{3}}{4} = \frac{2 + \sqrt{3}}{2}$$

$16 - 12 = 4, 8\sqrt{3} - 8\sqrt{3} = 0$


Exercise 1F

1 Simplify:

a $\frac{1}{\sqrt{5}}$

b $\frac{1}{\sqrt{11}}$

c $\frac{1}{\sqrt{2}}$

d $\frac{\sqrt{3}}{\sqrt{15}}$

e $\frac{\sqrt{12}}{\sqrt{48}}$

f $\frac{\sqrt{5}}{\sqrt{80}}$

g $\frac{\sqrt{12}}{\sqrt{156}}$

h $\frac{\sqrt{7}}{\sqrt{63}}$

2 Rationalise the denominators and simplify:

a $\frac{1}{1+\sqrt{3}}$

b $\frac{1}{2+\sqrt{5}}$

c $\frac{1}{3-\sqrt{7}}$

d $\frac{4}{3-\sqrt{5}}$

e $\frac{1}{\sqrt{5}-\sqrt{3}}$

f $\frac{3-\sqrt{2}}{4-\sqrt{5}}$

g $\frac{5}{2+\sqrt{5}}$

h $\frac{5\sqrt{2}}{\sqrt{8}-\sqrt{7}}$

i $\frac{11}{3+\sqrt{11}}$

j $\frac{\sqrt{3}-\sqrt{7}}{\sqrt{3}+\sqrt{7}}$

k $\frac{\sqrt{17}-\sqrt{11}}{\sqrt{17}+\sqrt{11}}$

l $\frac{\sqrt{41}+\sqrt{29}}{\sqrt{41}-\sqrt{29}}$

m $\frac{\sqrt{2}-\sqrt{3}}{\sqrt{3}-\sqrt{2}}$

3 Rationalise the denominators and simplify:

a $\frac{1}{(3-\sqrt{2})^2}$

b $\frac{1}{(2+\sqrt{5})^2}$

c $\frac{4}{(3-\sqrt{2})^2}$

d $\frac{3}{(5+\sqrt{2})^2}$

e $\frac{1}{(5+\sqrt{2})(3-\sqrt{2})}$

f $\frac{2}{(5-\sqrt{3})(2+\sqrt{3})}$

- E/P** 4 Simplify $\frac{3-2\sqrt{5}}{\sqrt{5}-1}$ giving your answer in the form $p+q\sqrt{5}$, where p and q are rational numbers. **(4 marks)**

Problem-solving

You can check that your answer is in the correct form by writing down the values of p and q and checking that they are rational numbers.


Mixed exercise 1

1 Simplify:

a $y^3 \times y^5$

b $3x^2 \times 2x^5$

c $(4x^2)^3 \div 2x^5$

d $4b^2 \times 3b^3 \times b^4$

2 Expand and simplify if possible:

a $(x+3)(x-5)$

b $(2x-7)(3x+1)$

c $(2x+5)(3x-y+2)$

3 Expand and simplify if possible:

a $x(x+4)(x-1)$

b $(x+2)(x-3)(x+7)$

c $(2x+3)(x-2)(3x-1)$

4 Expand the brackets:

a $3(5y+4)$

b $5x^2(3-5x+2x^2)$

c $5x(2x+3)-2x(1-3x)$

d $3x^2(1+3x)-2x(3x-2)$

5 Factorise these expressions completely:

a $3x^2 + 4x$

b $4y^2 + 10y$

c $x^2 + xy + xy^2$

d $8xy^2 + 10x^2y$

6 Factorise:

a $x^2 + 3x + 2$

b $3x^2 + 6x$

c $x^2 - 2x - 35$

d $2x^2 - x - 3$

e $5x^2 - 13x - 6$

f $6 - 5x - x^2$

7 Factorise:

a $2x^3 + 6x$

b $x^3 - 36x$

c $2x^3 + 7x^2 - 15x$

8 Simplify:

a $9x^3 \div 3x^{-3}$

b $(4^{\frac{3}{2}})^{\frac{1}{3}}$

c $3x^{-2} \times 2x^4$

d $3x^{\frac{1}{3}} \div 6x^{\frac{2}{3}}$

9 Evaluate:

a $\left(\frac{8}{27}\right)^{\frac{2}{3}}$

b $\left(\frac{225}{289}\right)^{\frac{3}{2}}$

10 Simplify:

a $\frac{3}{\sqrt{63}}$

b $\sqrt{20} + 2\sqrt{45} - \sqrt{80}$

11 a Find the value of $35x^2 + 2x - 48$ when $x = 25$.

b By factorising the expression, show that your answer to part a can be written as the product of two prime factors.

12 Expand and simplify if possible:

a $\sqrt{2}(3 + \sqrt{5})$

b $(2 - \sqrt{5})(5 + \sqrt{3})$

c $(6 - \sqrt{2})(4 - \sqrt{7})$

13 Rationalise the denominator and simplify:

a $\frac{1}{\sqrt{3}}$

b $\frac{1}{\sqrt{2} - 1}$

c $\frac{3}{\sqrt{3} - 2}$

d $\frac{\sqrt{23} - \sqrt{37}}{\sqrt{23} + \sqrt{37}}$

e $\frac{1}{(2 + \sqrt{3})^2}$

f $\frac{1}{(4 - \sqrt{7})^2}$

14 a Given that $x^3 - x^2 - 17x - 15 = (x + 3)(x^2 + bx + c)$, where b and c are constants, work out the values of b and c .

b Hence, fully factorise $x^3 - x^2 - 17x - 15$.

(E) 15 Given that $y = \frac{1}{64}x^3$ express each of the following in the form kx^n , where k and n are constants.

a $y^{\frac{1}{3}}$

(1 mark)

b $4y^{-1}$

(1 mark)

(E/P) 16 Show that $\frac{5}{\sqrt{75} - \sqrt{50}}$ can be written in the form $\sqrt{a} + \sqrt{b}$, where a and b are integers. **(5 marks)**

(E) 17 Expand and simplify $(\sqrt{11} - 5)(5 - \sqrt{11})$. **(2 marks)**

(E) 18 Factorise completely $x - 64x^3$. **(3 marks)**

(E/P) 19 Express 27^{2x+1} in the form 3^y , stating y in terms of x . **(2 marks)**

- (E/P)** 20 Solve the equation $8 + x\sqrt{12} = \frac{8x}{\sqrt{3}}$
Give your answer in the form $a\sqrt{b}$ where a and b are integers. (4 marks)
- (P)** 21 A rectangle has a length of $(1 + \sqrt{3})$ cm and area of $\sqrt{12}$ cm².
Calculate the width of the rectangle in cm.
Express your answer in the form $a + b\sqrt{3}$, where a and b are integers to be found.
- (E)** 22 Show that $\frac{(2 - \sqrt{x})^2}{\sqrt{x}}$ can be written as $4x^{-\frac{1}{2}} - 4 + x^{\frac{1}{2}}$. (2 marks)
- (E/P)** 23 Given that $243\sqrt{3} = 3^a$, find the value of a . (3 marks)
- (E/P)** 24 Given that $\frac{4x^3 + x^{\frac{5}{2}}}{\sqrt{x}}$ can be written in the form $4x^a + x^b$, write down the value of a and the value of b . (2 marks)

Challenge

- a** Simplify $(\sqrt{a} + \sqrt{b})(\sqrt{a} - \sqrt{b})$.
- b** Hence show that $\frac{1}{\sqrt{1} + \sqrt{2}} + \frac{1}{\sqrt{2} + \sqrt{3}} + \frac{1}{\sqrt{3} + \sqrt{4}} + \dots + \frac{1}{\sqrt{24} + \sqrt{25}} = 4$

Summary of key points

- You can use the laws of indices to simplify powers of the **same base**.
 - $a^m \times a^n = a^{m+n}$
 - $a^m \div a^n = a^{m-n}$
 - $(a^m)^n = a^{mn}$
 - $(ab)^n = a^n b^n$
- Factorising is the opposite of expanding brackets.
- A quadratic expression has the form $ax^2 + bx + c$ where a , b and c are real numbers and $a \neq 0$.
- $x^2 - y^2 = (x + y)(x - y)$
- You can use the laws of indices with any rational power.
 - $a^{\frac{1}{m}} = \sqrt[m]{a}$
 - $a^{\frac{n}{m}} = \sqrt[m]{a^n}$
 - $a^{-m} = \frac{1}{a^m}$
 - $a^0 = 1$
- You can manipulate surds using these rules:
 - $\sqrt{ab} = \sqrt{a} \times \sqrt{b}$
 - $\sqrt{\frac{a}{b}} = \frac{\sqrt{a}}{\sqrt{b}}$
- The rules to rationalise denominators are:
 - Fractions in the form $\frac{1}{\sqrt{a}}$, multiply the numerator and denominator by \sqrt{a} .
 - Fractions in the form $\frac{1}{a + \sqrt{b}}$, multiply the numerator and denominator by $a - \sqrt{b}$.
 - Fractions in the form $\frac{1}{a - \sqrt{b}}$, multiply the numerator and denominator by $a + \sqrt{b}$.